

TEP-11R  
June 1993  
hep-th/9303049

# Exact Solutions to the Wheeler-DeWitt Equation of Two Dimensional Dilaton Gravity

(revised)

**Takayuki Hori**<sup>†</sup>

*Institute for Physics, Teikyo University*

*Otsuka 359, Hachioji-shi, Tokyo 192-03, Japan*

## Abstract

The two dimensional dilaton gravity with the cosmological term and with an even number of matter fields minimally coupled to the gravity is considered. The exact solutions to the Wheeler-DeWitt equation are obtained in an explicit functional form, which contain an arbitrary holomorphic function of the matter fields.

---

<sup>†</sup>E-mail address: e00353@sinet.ad.jp

Recently the full quantum analysis of the two dimensional dilaton gravity has been considered to be of great importance because it must finally settle up the uncertainty on the black hole evaporation?? in two dimensions, the problem first posed by Callan et al.?? in a semi-classical approximation. There have been some attempts to analyse the model in the full quantum setting (see, *e.g.*, ref.(3)). Among them, however, the old-fashioned Wheeler-DeWitt (WD) quantization scheme?? has scarcely been given attention??. This may be partially due to the conceptual trouble one encounters in interpreting the wave function, when one is forced to rely on the asymptotic flatness to extract any measurable quantities. Moreover the exact solutions are hardly found except in some mini superspace models??.

In this note we find the latter problem, *i.e.*, to find the exact solutions to the WD equation is tractable in the two dimensional dilaton gravity. The problem on the interpretation of the wave function and the analysis of the physical outcomes are postponed for a future study.

The classical action?? is

$$I = \int d^2x e \{ e^{-2\phi} [R(\omega) + k(\nabla\phi)^2 + 4\lambda^2] - \frac{1}{2} \sum_a (\nabla f_a)^2 \}, \quad (1)$$

where  $\phi$  and  $f_a$  are the dilaton and matter fields, respectively, and  $\lambda$  is the cosmological constant.  $R(\omega)$  is the curvature expressed in terms of the spin connection  $\omega_{\mu a}{}^b$ , and  $e$  is the determinant of the zweibein,  $e_{\mu a}$ . The parameter  $k$  depends on models and  $k = 4$  corresponds to ref.(2), where the classical equations of motion are simplified. In the case  $k = 8$  the WD equations have a different type of solutions (see below).

The local Lorentz gauge is fixed by putting

$$e_{\mu a} = \begin{pmatrix} \alpha & \beta e^{-\rho} \\ 0 & e^\rho \end{pmatrix}, \quad (2)$$

where  $\alpha$  and  $\beta$  are the lapse function and the shift vector, respectively. In ref.(7) the WD equations without the cosmological term were obtained by the standard procedure??. For

the case with general  $k$  and with the cosmological term the WD equations are expressed as

$$\chi\Psi \equiv \left[ (\rho'(x) - \frac{\partial}{\partial x}) \frac{\delta}{\delta\rho(x)} + \phi'(x) \frac{\delta}{\delta\phi(x)} + \sum_a f'_a(x) \frac{\delta}{\delta f_a(x)} \right] \Psi = 0, \quad (3)$$

$$\begin{aligned} \Phi\Psi \equiv & \left[ \frac{1}{4} \frac{\delta^2}{\delta\rho(x)^2} + \frac{1}{k} \frac{\delta^2}{\delta\rho(x)\delta\phi(x)} - \frac{16}{k} e^{-4\phi(x)} \left( \frac{8-k}{4} \phi'^2 - \phi^{(2)} - \lambda^2 e^{2\rho} \right) \right. \\ & \left. + \frac{2}{k} e^{-2\phi(x)} \sum_a \left( \frac{\delta^2}{\delta f_a(x)^2} - f'_a(x)^2 \right) \right] \Psi = 0, \end{aligned} \quad (4)$$

where  $x = x^1$  and primes denote the derivatives with respect to  $x$ , and  $\phi^{(2)} \equiv \phi'' - \rho'\phi'$  is the second order covariant derivative in the spatial dimension. In eq.(4) we should regularize the functional derivations at the same points. The appropriate rule is determined by the requirement that the constraint algebra closes or by the consistent condition for the couple of the WD equations (3) and (4). This requires  $\frac{\delta\varphi(x)}{\delta\varphi(x)} = \delta(x, x) = 0$  for  $\varphi = (\rho, \phi, f_a)$ , and can be achieved by the dimensional regularization??.

To solve the WD equations let us assume the spatial manifold is compact and there are two matter fields,  $f_1$  and  $f_2$ . We make the following ansatz for the wave functional  $\Psi$

$$\Psi = \exp i \int dx [e^{-2\phi} \phi' g(X, \phi, z, \bar{z}) + \frac{1}{4} (z\bar{z}' - \bar{z}z')], \quad (5)$$

$$X \equiv e^{-\rho} \phi', \quad z \equiv f_1 + i f_2, \quad (6)$$

where  $g$  is a function to be determined. The  $\chi$ -constraint, eq.(3), amounts to the general coordinate invariance of the wave functional in the spatial dimension. Since  $\phi', z'$  and  $\bar{z}'$  are scalar densities and  $X, \phi, z$  and  $\bar{z}$  are scalars,  $\Psi$  satisfies the  $\chi$ -constraint. After some calculations we find the hamiltonian constraint, eq.(4), can also be satisfied for a suitable  $g$ . For  $k \neq 8$  the solution is

$$\Psi_{[A]} = \exp i \int dx \left[ -e^{-2\phi} \left( 2\phi'^{-1} \phi^{(2)} \Lambda + \frac{(e^{\frac{8-k}{2}\phi} A(z))'}{X^2(1+\Lambda)} \right) + \frac{1}{4} (z\bar{z}' - \bar{z}z') \right], \quad (7)$$

$$\Lambda \equiv \sqrt{1 - \left( \frac{4}{8-k} \lambda^2 + e^{\frac{8-k}{2}\phi} A(z) \right) X^{-2}}, \quad (8)$$

where  $A(z)$  is an arbitrary *holomorphic* function of  $z \equiv f_1 + i f_2$ . For  $k = 8$  we have a

slightly different form of the solution:

$$\Psi_{[B]} = \exp i \int dx \left[ -e^{-2\phi} \left( 2\phi'^{-1} \phi^{(2)} \tilde{\Lambda} + \frac{(B(z) - 2\lambda^2 \phi)'}{X^2(1 + \tilde{\Lambda})} \right) + \frac{1}{4}(z\bar{z}' - \bar{z}z') \right], \quad (9)$$

$$\tilde{\Lambda} \equiv \sqrt{1 - (B(z) - 2\lambda^2 \phi)X^{-2}}, \quad (10)$$

where  $B(z)$  is an arbitrary holomorphic function. It is straightforward to generalize the solution to the models with an even number of real matter fields.

If the cosmological term is absent there exists another type of solutions which cannot be expressed in the above form with  $\lambda = 0$ , *i.e.*,

$$\Psi_0 = \exp i \int dx [e^{-2\phi} (-2\phi'^{-1} \phi^{(2)} + M(\bar{z}, \phi)') + \frac{1}{4}(z\bar{z}' - \bar{z}z')], \quad (11)$$

where  $M$  is an arbitrary *anti*-holomorphic function of  $\bar{z}$  and  $\phi$ . The solution found in ref.(7) is a special case of eq.(11). Note that the space of the solutions expands at the point  $\lambda = 0$ . Although we work within the ansatz (5) this implies some sort of degeneracy at  $\lambda = 0$ .

Since the hamiltonian operator,  $\Phi$ , is real we see the complex conjugates of the above solutions are other independent solutions. Thus we have the most general solution

$$\Psi = \int \mathcal{D}A \left[ C_{[A]} \Psi_{[A]} + \tilde{C}_{[A^*]} \Psi_{[A^*]}^* \right], \quad (12)$$

where  $\Psi_{[A]}$  is one of the above solutions, and  $C_{[A]}$  and  $\tilde{C}_{[A^*]}$  are arbitrary coefficients depending on the function form of  $A$  and  $A^*$ . If one proceeds to the third quantization the coefficients  $C(\tilde{C})$  are interpreted as annihilation (creation) operators of the state specified by the particular configuration corresponding to the holomorphic function  $A(z)$ .

Our solutions must contain much informations on the quantum states of the two dimensional dilaton gravity, compared with those in the mini-superspace approaches, since they were obtained without any approximations. An investigation of the physical interpretation for the solution is expected to give a final answer to the problem of the black hole evaporation in two dimensions.

I wish to thank H.Kakuhata and M.Kamata for discussions. I am grateful to the members of High Energy Theory Group in Tokyo Metropolitan University for kind hospitality.

## References

- 1) S. W. Hawking, *Commun. Math. Phys.*, **43**, (1975) 199.
- 2) C. Callan, S. B. Giddings, J. A. Harvey and A. Strominger, *Phys. Rev.* **D45**, (1992) R1005.
- 3) S. Hirano, Y. Kazama and Y. Satoh, preprint UT-Komaba 93-3, Mar. 1993.
- 4) B. S. DeWitt, *Phys. Rev.* **160** (1967) 1113;  
J. A. Wheeler, in *Relativity groups and topology* (Gordon and Breach, New York, 1964).
- 5) A. Miković, Imperial-TP/92-93/08, Nov. 1992.
- 6) U. H. Danielsson, CERN-TH. 6711/92, Nov. 1992.
- 7) T. Hori and M. Kamata, preprint TEP-10, Feb. 1993.